

***Gnathia* sp. (Gnathiidae) Infestations on Marine Fish Species from Turkey**

Ali ALAŞ *  Ahmet ÖKTENER ** Muhittin YILMAZ ***

* Department of Science, Education Faculty, Aksaray University, 68100, Aksaray - TURKEY

** İstanbul Provincial Directorate of Agriculture, Directorate of Control, Kumkapı Fish Auction Hall, Aquaculture Office, Kumkapı, 34130 Istanbul - TURKEY

*** Department of Biology, Faculty of Science and Art, Kafkas University, 36100, Kars - TURKEY

Yayın Kodu (Article Code): 2008/96-A

Summary

This study was carried out between June and August 2006. Totally 150 samples out of ten different fish species caught from the Marmara Sea, the Black Sea and the Aegean Sea were examined. For this purpose, body surface, fins, gills and mouth cavity of each fish were examined with the aid of a dissecting microscope. After that, ectoparasite samples were preserved in 70% ethanol to be observed under the dissecting microscope for identification. *Gnathia* sp. was found out in the mouth cavity and gill filaments of these infected fishes. This is the first study reporting *Gnathia* sp. from *Serranus cabrilla*; *Mullus surmuletus*; *Mugil cephalus*; *Trachurus mediterraneus*; *Sarpa salpa*; *Sciaena umbra*.

Keywords: *Gnathia* sp., Sea of Marmara, Black Sea, Aegean Sea

Türkiye Deniz Balığı Türlerinde *Gnathia* sp. (Gnathiidae) İnfestasyonu

Özet

Bu çalışma Haziran-Ağustos 2006 periyodunda yapılmıştır. Marmara Denizi, Karadeniz ve Ege Denizi'nden yakalanan 10 farklı balık türünden toplam 150 balık incelenmiştir. Bu amaçla her bir balığın vücut yüzeyi, yüzgeçleri, solungaçları ve ağız boşluğu disseksiyon mikroskobu ile incelenmiştir. Bundan sonra ektoparazit örnekleri disseksiyon mikroskobu altında incelenmiş olup, toplanan balıkların teşhis edileceği zamana kadar %70'lik etanolde muhafaza edilmiştir. *Gnathia* sp. larvalarına enfekte balıkların solungaç filamentleri ve ağız boşluklarında rastlanmıştır. Bu çalışmayla Türkiye'de *Serranus cabrilla*; *Mullus surmuletus*; *Mugil cephalus*; *Trachurus mediterraneus*; *Sarpa salpa*; *Sciaena umbra* balık türlerinden *Gnathia* sp. ilk defa rapor edilmektedir.

Anahtar sözcükler: *Gnathia* sp., Marmara Denizi, Karadeniz, Ege Denizi

INTRODUCTION

Gnathiid isopods (Crustacea: Isopoda: Gnathiidae) have complex life cycles with free-living adults and three stages of parasitic juveniles that feed on the blood and tissue fluids of teleosts and elasmobranchs. Juveniles have piercing and serrated mouthparts, including paired toothed mandibles and maxillules (maxillae 1), grooved paragnaths (hypopharynx), robust maxillipeds and hooked gnathopods. Young gnathiids can thus attach and feed on the

surfaces of fish, sometimes in large numbers. Whereas gnathiid isopods as larvae are external fish parasites, as adults they live in cavities and crevices on the sea floor, such as in sponges, sea anemones and tunicates, coral rubble, or sediment cavities ¹⁻⁴. Presently there are 182 identified species of gnathiids reported from the world oceans ⁵. Larvae, adult females and males differ considerably in their shape and behaviour. Praniza which is parasitic larval form of *Gnathia*



İletişim (Correspondence)



90 382 2801146 & +90 505 8435643



alasali@hotmail.com

sp. ⁶. Praniza feed on their hosts' blood. There is enormous expansion of the mid-gut to accomodate these blood meals. When the ectoparasitic phase is over, the praniza leave their host and moult into the adult male or female. The adults are cryptic, do not feed and lead secluded lives in sheltered places on the sea-bed ^{7,8}. Numerous species have been reported in fish culture including *Gnathia vorax* on Sparidae, Serranidae, Mugilidae from the Mediterranean coast of Israel (9); *Gnathia* sp. on salmon farm (10); Paragnathia sp. on *Anguilla anguilla* ¹¹; *Gnathia* sp. on atlantic salmon ¹².

This study is specifically aimed to investigate the infestations and hosts distribution of *Gnathia* sp. from Turkey.

MATERIAL and METHODS

A total of 150 fish belonging to several family were caught from three sea as follows: *Scorpaena scrofa*, *Serranus cabrilla*, and *Pagellus erythrinus* from the Sea of Marmara; *Mullus surmuletus* from Black Sea; *Mugil cephalus*, *Gaidropsarus mediterraneus*, *Scorpaena scrofa*, *Trachurus mediterraneus*, *Sarpa salpa*, *Serranus cabrilla*, *Diplodus vulgaris*, and *Sciaena umbra* from Aegean Sea. They were examined for ectoparasites in July-August 2006. After capturing, the fishes were placed on ice for approximately 6 hr and the body surface, fins, gills, mouth of each fishes were examined. The gill arches were preserved in 70%

ethanol and later examined with the aid of a dissecting microscope (American Optycal Microscope). The parasites were removed and preserved in 70% ethanol. The identification of parasites were completed according to various authors ^{6-8,13}.

RESULTS

Infestation values: Number of fish examined, number of fish infested, prevalence (%), intensity (min-max), total number of parasites were given in Table 1.

Table 1. Parasitological index of *Gnathia* sp. collected from the seas surrounding Turkey

Tablo 1. Türkiye'yi çevreleyen denizlerden toplanan *Gnathia* sp.'nin parazitolojik indexi

Fish Species	NFE	NFI	P	MI (min-max)	NP
<i>Mullus surmuletus</i>	23	2	8.7	1.5(1-2)	3
<i>Scorpaena scrofa</i>	26	3	11.5	3.3(1-6)	10
<i>Serranus cabrilla</i>	34	5	14.7	2.6(1-3)	13
<i>Mugil cephalus</i>	2	2	100	10(5-15)	20
<i>Gaidropsarus mediterraneus</i>	8	4	50	3.5(2-6)	14
<i>Trachurus mediterraneus</i>	5	2	40	2.0(1-3)	4
<i>Sarpa salpa</i>	4	3	75	2.6(1-4)	8
<i>Diplodus vulgaris</i>	15	6	40	3.6(3-8)	22
<i>Sciaena umbra</i>	6	6	100	4.6(4-8)	28
<i>Pagellus erythrinus</i>	27	7	26	3.0(1-5)	21

NFE = Number of fish examined; **NFI** = Number of fish infested; **P** = Prevalence (%); **MI (min-max)** = Mean intensity (minimum-maximum); **NP** = Total number of parasites




Fig 1. *Gnathia* sp.

Şekil 1. *Gnathia* sp.

DISCUSSION

There are various studies about *Gnathia* sp. in the Black Sea and Mediterranean Sea. *Gnathia oxyure* and *Gnathia bacescoi* species from the Black Sea were reported by Kononenko ¹⁴. In addition, Gnathiids in the Mediterranean Sea were investigated by Monad ¹.

In a research done by Papoutsoglou, *Gnathia maxillaris* was reported on several fishes from Greece: *Apogon imberbis*, *Boops salpa*, *Cantharus lineatus*, *Chromis chromis*, *Coris julis*, *Crenilabrus mediterraneus*, *C. quinque maculatus*, *C. scina*, *C. tinca*, *Diplodus annularis*, *D. vulgaris*, *Gobius jozo*, *G. ophiocephalus*, *Labrus merula*, *Maena maena*, *Mullus barbatus*, *M. surmuletus*, *Onos tricirratu*, *Pagellus erythrinus*, *Phycis blennioides*, *Puntazzo puntazzo*, *Scorpaena porcus*, *Serranus scriba*, *Sphyaena sphyaena*, *Thalassoma pavo*, *Trachinus raco*, *Trigla lineata*, *Umbrina cirrosa*, *Uranoscopus scaber*, and *Zeus faber* ¹⁵. At Adriatic Sea, *Gnathia* sp. (*vorax*) from *Merluccius merluccius*, *Pagellus erythrinus*, *Mullus barbatus*, *Trigla lyra*, and *Gnathia* sp. (*piscivora*) from *Chelon labrosus* were identified by Radujkovic ¹⁶.

In another study, *Gnathia* sp. was reported from *Uranoscopus scaber*, *Serranus scriba*, *Onos tricirrata*, *Acipenser stellatus*, *Ophidium barbatum*, *Trigla hirundo*, and *Crenilabrus pavo* from Black Sea ¹⁷.

The Gnathiids previously from Turkish sea are reported as both free and parasitic phase. *Gnathia vorax* was determined by some authors in holes and in wooden boards from Black Sea and Aegean Sea as a free form ^{18,19}. *Gnathia* sp. praniza larvae were reported from several fishes from Aegean Sea and Mediterranean Sea of Turkey ¹⁹⁻²³. *Gnathia* sp. was noted on *Diplodus annularis*, *D. vulgaris*, *Spondyliosoma cantharus*, *Lithognathus mormyrus*, *Diplodus sargus*, *Dentex dentex*, *Sparus pagrus*, *Crenilabrus tinca*, *Scorpaena porcus*, *S. scrofa*, *Gaidropsarus mediterraneus*, and *Umbrina cirrosa* from Gökçeada by Akmirza ^{21,23}, and *Gnathia* sp. on *Epinephelus aeneus* was found by Genç et al. ²⁰.

Mugil cephalus usually in schools over sand or mud bottom, *Sciaena umbra* occurs in shallow coastal waters mainly on rocky and sandy bottoms, *Sarpa salpa* found over rocky substrates and sandy area with algal growth, *Gaidropsarus*

mediterraneus lives generally at shallow depths near the shores on rocky bottom with aquatic vegetation ²⁴. Prevalance values of *Mugil cephalus*, *Sciaena umbra*, *Sarpa salpa* and *Gaidropsarus mediterraneus* were occurred over 50% (Table 1). These high prevalence values is likely to be related to their habitat since adults and larvae of gnathiids are located on the sea bed, these fish are probably more easily infected than the others.

Gnathia spp. are *protandrous* hermaphrodites and to form male individuals, culture medium is necessary. We were not able to prepare culture medium. So, we could not culture pranizas. For this reason, we were able to make the identification of this species at genus level.

Gnathia species were found belonging to four different families which are recorded in Turkey for first time. In this sampling, two main body areas were infested by *Gnathia* sp. in Turkish fishes; the mouth base and the gill filaments. The study shows that five demersal, four benthopelagic and one pelagic fish were infested with *Gnathia* sp.

ACKNOWLEDGEMENT

We are grateful to Dr. N.J. Smit from Department of Zoology, University of Johannesburg, South Africa for verifying the identity of *Gnathia* spp.

REFERENCES

1. Monod T: Les Gnathiidae. Essai monographique (Morphologie biologie systematique). Mem Soc Sci Nat Maroc, 13.1-668. 1926.
2. Upton NPD: Asynchronous male and female life cycles in the sexually dimorphic, harem-forming isopod *Paragnathia formica* (Crustacea: Isopoda). J Zool, 212, 677-690, 1987.
3. Klitgaard AB: The distribution and habitats in the North Atlantic of two gnathiid species (Crustacea, Isopoda) and their reproductive biology in the Denmark Strait and north of Iceland. Meddelelser om Grønland, Biosci, 47, 1-32, 1997.
4. Smit NJ, Davies A J: The curious life-style of the parasitic stages of gnathiid isopods. Adv Parasit, 58, 289-391, 2004.
5. Kensley B, Schotte M, Schilling S: World list of marine, freshwater and terrestrial isopod crustaceans. Smithsonian Institution, Washington, DC, USA. Available from <http://www.nmnh.si.edu/iz/isopod/>. Accessed: 10 May 2008. 2005.
6. Möller H, Anders K: Diseases and Parasites of Marine Fishes. Verlag Heino Möller, Kiel, 1986.
7. Kabata Z: Diseases of Fishes: Book I. Crustaceans Enemies of Fishes. T.F.H. Publications at New Jersey US, 1970.
8. Naylor E: British Marine Isopods. Academic Pres at New York, 1972.

- 9. Paperna I, Por FD:** Preliminary data on the Gnathiidae (Isopoda) of the Northern Red Sea, the Bitter Lakes, and the Mediterranean and the biology of *Gnathia piscivora* n. sp. Rapp P-v Reün Commn int *Explor Mer MeÄditerr*, 24, 195-197, 1977.
- 10. Palmer MA:** Invertebrate drift: behavioral experiments with intertidal meiobenthos. *Mar Behav Phy*, 10, 235-253, 1984.
- 11. Mugridge RER, Stallybrass HG:** A mortality of eels (*Anguilla anguilla* L.) attributed to Gnathiidae. *J Fish Biol*, 6, 81-82, 1983.
- 12. Drinan EM, Rodger HD:** An occurrence of *Gnathia* sp. ectoparasitic isopods, on caged Atlantic salmon. *Bull Eur Ass Fish Path*, 10, 141-142, 1990.
- 13. Smit NJ, Van As JG, Basson L:** A redescription of the adult male and larvae of *Gnathia africana* Barnard, 1914 (Gnathiidae: Crustacea: Isopoda) from Southern Africa. *Folia Parasit*, 46, 229-240, 1999.
- 14. Kononenko AF:** Parasitic isopoda of the Black Sea fishes. V. Symposium on Medical and Vet Acaroent, Gdansk, 1985.
- 15. Papoutsoglou SE:** Metazoon parasites of fishes from Saronicos Gulf Athens-Greece. *Thalassog*, 1 (1): 69-102, 1976.
- 16. Radujkovic BM:** Isopoda-parasites of the South Adriatic economically important fish species. *Acta Adriat*, 23 (1/2): 153-161, 1982.
- 17. Reshetnikova AV:** Parazitofauna nekotorykh promislovykh ryb *Chernogo morya*. Autoreferat Dissertation, Institut Zoologii, Akademia Nauk SSSR, 1954.
- 18. Gönlügür G:** The qualitative and quantitative investigations on some facies of the upper-infralittoral zone of the Western black sea (Sinop) shores. Ege University, *PhD Thesis*. Science Institution, İzmir, 2003.
- 19. Geldiay R, Kocataş A:** Isopods collected in İzmir bay, Aegean Sea. *Crustaceana, Suppl. 3, Studies on Peracarida*, 19-30, 1972.
- 20. Genc M, Genc A, Can MF, Genc E, Cengizler I:** A first documented record of Gnathiid infestation on white grouper (*Epinephelus aeneus*) in Iskenderun Bay (Northeastern Mediterranean), Turkey. *J Appl Ichthyol*, 21 (5): 448-450, 2005.
- 21. Akmirza A:** The samples from metazoon parasites detected in fish around Gökçeada. *Congress of National Aegean Islands*, 2001.
- 22. Kırkım F:** Investigations on Systematic and Ecology of the Eagean Sea Isopoda Fauna. Ege University, PhD Thesis. Science Institution, İzmir, 1998.
- 23. Akmirza A:** Seasonal distribution of parasites detected in fish belonging to the Sparidae family found near Gökçeada. *J Parasit Turkey*, 24 (1): 435-441, 2000.
- 24. Froese R, Pauly D:** Fish base. World wide web electronic publication. www.fishbase.org. Accessed: 5 June 2008. 2008.